

Bionet-Winterweizenversuche Ost (Niederösterreich)

Standortübersicht:

Standort:	Pellendorf			Maissau			Langau		
Vorfrucht:	Luzerne/Rotklee			Ölkürbis			Luzerne		
Boden	Ertrag	Protein	Roherlös	Ertrag	Protein	Roherlös	Ertrag	Protein	Roherlös
	kg/ha, relativ	%	€/ha, relativ	kg/ha, relativ	%	€/ha, relativ	kg/ha, relativ	%	€/ha, relativ
Albertus	99%	14,3%	102%				81%	13,7%	90%
Alessio				97%	10,6%	97%			
Alicantus									
Arminius	111%	14,5%	115%	91%	10,5%	91%	99%	14,6%	113%
Arnold	104%	15,0%	107%	91%	11,7%	114%	90%	14,8%	102%
Aurelius									
Bernstein									
Edelmann	101%	14,5%	104%	79%	11,7%	99%	99%	13,1%	110%
Ehogold	102%	14,0%	105%				91%	13,5%	101%
Element				100%	10,4%	100%			
Lukullus				110%	10,1%	110%			
Tilliko	88%	14,7%	90%	96%	9,9%	96%	86%	13,4%	96%
Tobias	104%	14,7%	107%	90%	11,1%	113%	94%	15,0%	107%
Capo	2.992	13,8%	1.280	3.394	10,0%	950	5.318	12,7%	2.047
Stabw Capo	5%			4%			1%		

Standort:	Absdorf			Ebergassing			Pöchlarn		
Vorfrucht:	Luzerne			Mais			Soja		
Boden	Ertrag	Protein	Roherlös	Ertrag	Protein	Roherlös	Ertrag	Protein	Roherlös
	kg/ha, relativ	%	€/ha, relativ	kg/ha, relativ	%	€/ha, relativ	kg/ha, relativ	%	€/ha, relativ
Albertus									
Alessio	89%	14,9%	89%						
Alicantus				112%	11,5%	112%	86%	12,6%	94%
Arminius	92%	15,0%	92%						
Arnold				93%	12,3%	102%	84%	13,3%	103%
Aurelius				94%	12,1%	103%	101%	12,4%	111%
Bernstein	88%	14,0%	88%						
Edelmann	98%	13,4%	95%	86%	11,1%	86%	94%	11,7%	94%
Ehogold	101%	14,4%	101%						
Element									
Lukullus									
Tilliko	88%	14,8%	88%	52%	13,5%	64%	87%	12,1%	96%
Tobias									
Capo	5.261	14,0%	2.315	2.987	11,2%	1.051	4.656	11,9%	1.639
Stabw Capo				9%			3%		

Stabw Capo ... die Standardabweichung ist ein Maß für die Streuung der Ertragswerte des jeweils dreifach wiederholten Standards Capo an den einzelnen Versuchsstandorten.